


## APPENDIX 7 GIS DATA: ASSUMPTIONS AND LIMITATIONS

In order to carry out the study, a review of provided spatial data was carried out in order to assess its quality for purpose. Where data gaps were identified, additional data was requested (where available) through the appropriate LGAs, and/or supplemented from SMEC own internal data holdings.

### Assumptions

The data review highlighted various issues with the supplied data, that needed to be addressed prior to desktop analysis, to ensure that results produced would be valid. Of highlight:

- The waterline from the DECC provided (sourced from the DTDB) Hydrographic dataset is the did not accurately correspond to the HWM in some areas, and in a small amount of areas, was outside of the watercourse as viewed on current aerial imagery. This is seen below with the supposed DTDB waterline in red, and the SMEC-adjusted waterline, matching the existing feature, in blue.



If not adjusted, this issue would have led to error in vegetation and inundation calculation. SMEC reviewed the entire waterline for the study area to the provided aerial imagery, and amended it where required, and used this as the basis for the HWM for the study.

- Differences in LGA boundaries were evident between different sets from different sources. All define the relevant boundaries, but the issue is relevant in this study as each source set of LGA boundaries from different agencies (LGAs, Dept. Lands, ABS, etc), has a different definition of how far into the Georges River Estuary the LGA boundary extends; this is visualised below. Here, the DTDB-sourced LGA boundaries as provided by DECC are shown in blue, and extend to the centre of the watercourse. The ABS LGA boundaries on the other hand (in red), do not extend into the watercourse. There also minor differences in boundary, as seen in the image. These differences would lead to differences in calculations for vegetation coverage of an LGA, and especially for Seagrass calculations, within the main water channel.



SMEC made the decision to go with the provided DTDB boundaries, to ensure consistency. As the boundary is represented as the centre of the watercourse, this obviously needs to be considered as somewhat arbitrary when considering channel-based vegetation as Seagrass divided by this line.

### **Sea Level Rise**

Contours for the estimation of impact due to sea level change were calculated using elevation data as provided by the various councils. There was no uniform-scale elevation data provided by all authorities, with intervals ranging from 50cm to 10m between contours. This level of difference meant that creation of an elevation model across the whole study area would be subject to varying accuracies, but still suitable for indicative estimation of areas of potential future inundation.

To generate the elevation model for the whole study area, the following methodology was used. All provided elevation data was combined (with additional elevation data from SMEC's internal holdings, where authorities were unable to provide any coverage), into a single set. This data was then used as the input to create a Digital Elevation Model (DEM) of the study area using FME (Safe Software 2009). By combining all available elevation data into the modelling process rather than producing one DEM per authority, it was possible to use neighbouring authorities data to allow the DEM creation process to create a more accurate

model. Once the DEM was created, contours were taken from this at 50cm and 90cm, the estimated sea level rise for 2050 and 2100 respectively. It should be noted that in areas where available contour data was limited (i.e. only 10m intervals), anomalies can appear due to fine scale of the required contours in the output vs the inputs. In this instance, such areas noted as subject to inundation should be treated as indicative only, and further study on sea level change for the Georges River estuary should be conducted using high resolution elevation data set that covers the whole of catchment, such as LiDAR. It should also be noted that sea level changes are also subject to flood protection works, urban development etc, which is not captured in the provided contour data, which provides a 'bare-earth' model of the ground. Using LiDAR for future study would avoid this issue, as it would capture all above ground infrastructure in the study area, allowing for more accurate inundation area modelling and offsetting.

### **Precision of Vegetation Mapping (DECCW, 2009)**

This layer contains digital mapping of the vegetation communities of the Sydney Metropolitan Catchment Management Authority area. Derived from Aerial Photographic Interpretation with communities defined using survey and statistical analysis. The layer includes the boundaries of vegetation communities and lists attributes including vegetation community, understorey, disturbance type & severity as described in the report: DECCW (2009) The Native Vegetation of the Sydney Metropolitan Catchment Management Authority Area. Unpublished report funded by the Australian Government and the Sydney Metro Catchment Management Authority. Department of Environment, Climate Change & Water, Hurstville.

Aerial photo interpretation was carried out between 1:1000 and 1:2000. The minimum polygon size was approximately 25m<sup>2</sup>, however some smaller polygons were interpreted at the interpreters discretion.

Linework was derived from 10cm resolution digital RGB orthographic aerial photography flown in 2005 & 2007 by Sinclair Knight Mertz (provided by SMCMA), using Definiens Developer 7 (refer to Vol 1 for more details). For the Holsworthy Defence area, 2008 50cm orthorectified digital imagery provided by the Department of Lands was used. Linework was cleaned and edited manually using ArcGIS 9.2 at a viewing scale of 1:2000.

Precision is 10m to 100m Determination: Deductive estimate. Ad-hoc comparisons with 1:25,000 layer data and various sorts of imagery indicated good correlation in terms of shape and size but with errors of the order indicated above in terms of position.

### **Seagrass calculations for unincorporated data**

Boundaries of LGAs- Rockdale boundary doesn't extend into Botany Bay whereas Sutherland does. Therefore 'Unincorporated' seagrass area in North Botany Bay is the area of Botany Bay adjacent to Rockdale council but not including Sutherland's area.